

Photochemical Nozaki–Hiyama–Kishi Coupling Enabled by Excited Hantzsch Ester

Yonghong Liu,[§] Shuangjie Lin,[§] Dandan Zhang, Bingkun Song, Yunhe Jin, Erjun Hao, and Lei Shi^{*†}



Cite This: *Org. Lett.* 2022, 24, 3331–3336



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: This work reports the first photochemical Nozaki–Hiyama–Kishi coupling enabled by bioinspired Hantzsch ester. The salient feature of this process is that commercially available and low-cost organic photoactive Hantzsch ester can serve as both an electron and a proton donor to reduce Cr/Ni to low-valent species and hydrolyze the Cr^{III}-alkoxy bond, thus bypassing the use of stoichiometric metallic reductants and additives such as TMSCl and Cp₂ZrCl₂. The mild conditions and operationally easy method showed broad compatibility with various alkenyl triflates and aldehydes, including electron-poor pentafluorobenzaldehyde which failed under previous conditions.




Chromium-mediated carbon–carbon forming reactions played a key role in contemporary organic synthesis.^{1–5} Particularly, the addition of vinyl-organochromium reagents to carbonyls, known as Nozaki–Hiyama–Kishi (NHK) coupling, has been proven to be a powerful and efficient method to construct valuable allylic alcohols. Because of unique features, such as excellent chemoselectivity and broad compatibility with functional groups, NHK reactions have been widely used in medicinal chemistry and total synthesis of natural products.⁴ Initially, NHK coupling required a Ni catalyst and a stoichiometric amount of CrCl₂ (Figure 1a).^{6–9} Later, a catalytic protocol was developed using Mn(0) as terminal reductants, thereby achieving the asymmetric induction with a chiral ligand.^{3,10–13} Alternatively, an electrochemical NHK reaction was reported bypassing the use of superstoichiometric metallic reducing agents (Figure 1b).^{14–16} In terms of reaction

mechanism, oxophilic additives such as TMSCl or Cp₂ZrCl₂ are indispensable for the NHK reaction, as they play the role of cleaving the Cr^{III}-alkoxy bond and releasing free Cr^{III} species in the catalytic cycle. However, these Lewis acid type additives can also activate aldehydes through coordination with the oxygen atom. Especially for electron-poor aryl-aldehydes, this activation led to single-electron reduction of aldehydes by low valence CrCl₂ producing undesired pinacols via ketyl radicals.¹⁰ Additional operation of aqueous workup in the presence of *n*-Bu₄NF to remove the silyl group of the crude product is required. Given the prevalent utilization of NHK couplings in the synthetic and industrial community, the development of sustainable conditions to avoid the use of superstoichiometric metallic reductants and exogenous additives and further expand the substrate scope is still highly desired.

In recent years, the merger of photoredox catalysis and transition metal catalysis, termed metallaphotoredox catalysis, has become a powerful and enabling tool in organic synthesis, where light-absorbing metal-polypyridyl complexes or organic dyes can modify the oxidation state of metal catalysts.^{17–20} In this regard, the first report of dual chromium and photoredox catalysis for Nozaki–Hiyama allylation of aldehydes was from the Glorius group in 2018.^{21,22} Later, Kanai and co-workers developed an asymmetric allylation by judicious choice of bisoxazoline ligand providing rapid access to enantiomerically pure homoallylic alcohols from simple unactivated alkenes.²³


various strategies for Nozaki–Hiyama–Kishi reaction



a) chemical strategy: stoichiometric Mn / additive: Me₃SiCl

b) electrochemical strategy: (+)Al/(-)Ni foam / additive: Cp₂ZrCl₂

c) photochemical strategy (this work)



- ◆ first photochemical Nozaki-Hiyama-Kishi reaction
- ◆ HE* as photoreductant
- ◆ free of additives and metallic reductants

Figure 1. State of the art of NHK reaction.

Received: March 12, 2022

Published: April 12, 2022



However, photo-promoted Cr/Ni catalyzed NHK alkenylation of aldehydes producing allylic alcohols remains underdeveloped.

Living organisms and plants sophisticatedly use photoactive cofactors, such as NADH and flavin, to convert light energy into chemical energy to fuel metabolic activities.^{24–27} The analog of NADH, 1,4-dihydropyridine (Hantzsch ester, HE), is well understood as a hydride source in the ground state and sacrificial electron donor to the photocatalyst in photochemical reactions.²⁸ Very recently, utilization of photoexcited HE* as a potential photoreductant (−2.28 V versus SCE) for metal-laphotoredox catalysis appealed to great interests. Melchiorre and Molander respectively reported that photoexcited HE* and its analogs can reduce Ni to low valence catalytically active species for C–C bond coupling.^{29,30} Inspired by these achievements, we hope to harness the reduction potential of HE* for chromium catalysis.³¹ Herein, we report the first photochemical NHK coupling of alkenyl triflates and aldehydes enabled by photoexcited HE* (Figure 1c).

The multiple roles of HE are to reduce Cr and Ni to active low-valent species and serve as proton sources to break the Cr^{III}-alkoxy bond, thus bypassing the use of stoichiometric metallic reductants and additives such as TMSCl or Cp₂ZrCl₂. Another advantage of photochemical NHK is that easily single-electron reduced substrates are accommodated and produced desired allylic alcohols when classical conditions failed.

The study was started by examining phenylpropionaldehyde **1a** and vinyl-triflate **2a** derived from 4-oxacyclohexanone as coupling partners. After variation of different reaction parameters, we found the suitable conditions in which a mixture of **1a** (0.5 equiv), **2a** (1 equiv), CrCl₂ (20 mol %), NiBr₂ (2 mol %), 4,7-Diphenyl-1,10-phenanthroline L₃ (2.2 mol %), and HE (3.0 equiv) in acetonitrile with irradiation of a 400 nm light emitting diode (LED) lamp for 48 h afforded the desired product **5a** with a satisfactory yield of 76% (Table 1, entry 1). Ligands are found to be important for the high yield of product.³² In the absence of any ligand, the yield of **5a** was only 43% (entry 1). When 1,10-Phenanthroline L₁ and Neocuproine L₂ were used the yields increased slightly (entries 3 and 4). Reactions were also sensitive to the light source. For example, irradiation with white LED, 450 nm LED, or 530 nm LED light, instead of 400 nm LED light, all resulted in very low yields of product **5a** (entries 6 and 7). This indicated that 400-nm-LED light is essential for the excitation of HE. Bench stable CrCl₃ was found to be suitable for the reaction affording the product **5a** in a slightly lower yield of 65% (entry 8). Several photocatalysts were tested for the reaction. However, in all cases the yields were relatively low and the conversion of **2a** was incomplete (entries 9–11). While additives such TMSCl and Cp₂ZrCl₂ were necessary for the classical catalytical approaches, it was not the case for the photoconditions and led to the decomposition of substrates (entries 12 and 13). Various solvents, including DMF, DMSO, and acetone, were evaluated, and they all resulted in less yields (entries 14–16). When reactions were performed in THF and DCM, no obvious product was formed. Moreover, the reaction did not proceed in the absence of HE, Cr, Ni, or light (entry 17).

With the optimal conditions in hand, we further investigated the scope of substrates. Remarkably, the reaction went well with various aliphatic/aromatic aldehydes **1** and alkenyl triflates **2**, affording the allylic alcohols in generally good yields, as depicted in Scheme 1. First, a broad range of functionalized aliphatic aldehydes were examined, including

Table 1. Reaction Optimization^{a,b}




entry	variation from standard conditions	yield (%)
1	None	80 (76) ^c
2	No ligands	43
3	L ₁ instead of L ₃	62
4	L ₂ instead of L ₃	58
5	L ₄ instead of L ₃	70
6	white LED instead of 400 nm LED	<5%
7	450 nm, or 530 nm LED, instead of 400 nm LED	<5%
8	CrCl ₃ instead of CrCl ₂	65
9	4CzIPN (450 nm LED)	38
10	Ir-1 (450 nm LED)	51
11	Ir-2 (450 nm LED)	42
12	TMSCl (1 equiv) added	18
13	Cp ₂ ZrCl ₂ (1 equiv) added	31
14	DMF instead of CH ₃ CN	47
15	DMSO instead of CH ₃ CN	38
16	Acetone instead of CH ₃ CN	61
17	No light/HE/CrCl ₃ /NiBr ₂	0

^aReaction conditions: **1a** (0.5 mmol), **2a** (1 mmol). ^bYields were determined by ¹H NMR spectroscopy versus an internal standard. ^cIsolated yield (details see Supporting Information (SI) S4).

primary aldehydes **3f–3g** and secondary aldehydes **3a–3b**, yielding allyl alcohols exclusively in comparable yields. Heteroatom substituents at the terminus of alkyl chains, such as TBS protected oxygen **3d**, phthalimide **3j**, Boc-protected amine **3e**, and even chlorine atom **3c**, are accommodated perfectly. Then the scope with respect to aromatic aldehydes was investigated under the standard conditions.

It is noteworthy that, in the classical metal-reduction conditions, for the electron-poor aromatic aldehydes the yields of NHK reaction are usually extremely low, accompanied by the production of pinnacle byproducts. Although it reported that desired allylic alcohols could be obtained by lowering the temperature to 0 °C, it is limited to monosubstituted chlorine or bromine atoms. Fortunately, under the photoreductive conditions, both substrates with electron-rich groups (**4b**, **4i**) and electron-poor groups, such as F- **4k**, CF₃- **4l**, CN- **4m**, CO₂Me- **4n**, and even pentafluorobenzaldehyde **4o**, were accommodated well. For the comparison with the metal reduction conditions, these electron-poor aromatic aldehydes were resubjected to the reaction in the presence of Mn and TMSCl, but no obvious alcohol products were observed (**4k–4o** yields shown in parentheses). Various alkenyl triflates were investigated, and the results showed that regarding functionalization various structure complicated aldehydes derived from (**5f**) were well tolerated. To further illustrate the potential applicability of this protocol in the late-stage heterocycloalkenyl triflates (**5a**, **5b**), spiro cycloalkenyl triflate (**5c**), cyclic alkenyl triflate (**5d**, **5e**), and linear alkenyl triflate natural products and active drugs were examined, including probenecid, oxaprozin, indomethacin, gemfibrozil, and bezafibrate, which all proceeded well under these conditions, producing the corresponding products in comparable yields (**6a–6e**). Overall, the NHK reactions of aldehydes and alkenyl triflates were carried out quite efficiently under photo-

Scheme 1. Scope of Photochemical NHK Reaction^{a,b,c}

^aReactions were performed on a scale of **1** (0.50 mmol, 1.0 equiv), **2** (1.00 mmol, 2.0 equiv), NiBr₂ (2 mol %), L₃ (2.2 mol %), CrCl₂ (20 mol %), HE (3.0 equiv), 400 nm LEDs, CH₃CN (0.1 M), isolated yield. ^bWhen the products and Hantzsch pyridine cannot be completely separated, yields were determined by ¹H NMR spectroscopy versus an internal standard. ^cYields in parentheses based on catalytic metal-reduction conditions: **1** (0.50 mmol, 1.0 equiv), **2** (1.00 mmol, 2.0 equiv), NiCl₂ (3 mol %), CrCl₂ (15 mol %), Mn (1.68 equiv), TMSCl (2.4 equiv), DME: DMF = 20:3 (0.1 M), 50 °C, 24 h [ref 10].

conditions without the use of exogenous additives and metallic reductants.

The redox potential of HE* was estimated to be −2.28 V versus SCE,³³ and thus the single electron reduction of Cr^{III} ($E_p\text{Cr}^{\text{III}/\text{II}} = -0.51$ V vs SCE in DMF)³⁴ and Ni^{II} ($E_{p/2}\text{Ni}^{\text{II}/\text{I}} = -1.2$ V vs SCE in DMF)³⁵ by HE* should be thermodynamically feasible. However, the excited-state lifetime of HE* is short ($\tau_0 = 320$ ps),³³ and it is not clear whether metals can be

efficiently reduced by HE* through the SET process. Therefore, a series of experiments and Stern–Volmer analysis were performed. The mixture of CrCl₃ (purplish-red solid) and HE in DMF was irradiated with a 400 nm LED at room temperature. The color of DMF changing to green indicated that CrCl₂ species was formed. The results of analysis by ultraviolet–visible (UV–vis) spectroscopy further revealed that the typical peak of CrCl₂ at 670, 652, and 476 nm

gradually appeared within 15 min, suggesting that the photoreduction process is fast (Figure 2a). On the contrary,




Figure 2. (a) UV-vis spectra showing the reduction of Cr^{III} to Cr^{II} by excited HE* upon photoirradiation with a 400 nm LED. (b) UV-vis spectra of Cr^{II}. (c) HE emission quenching by CrCl₃(THF)₃. (d) HE emission quenching by NiCl₂(DME) (for details, see SI S6).

when the same reaction mixture was exposed to irradiation with a 450 nm LED, the reduction is extremely slow (over 12 h). This finding indicated that using 400 nm light is crucial for the excitation of HE. Importantly, no change of color was observed in the dark even at elevated temperatures, indicating that ground-state SET was not occurring. Then fluorescence quenching experiments were performed to investigate the electron transfer in the SET process (Figure 2c, 2d). The luminescence of HE at its λ_{max} (438 nm) in degassed THF was quenched rapidly by both CrCl₃(THF)₃ and Ni^{II} (SI S6), supporting the notion that HE could act as an efficient photoreductant. The UV-visible absorption spectrum of two mixtures of HE with Cr and with Ni were identical with their individual spectra, indicating the lack of charge-transfer interactions in the ground state (SI S5). Therefore, fluorescence quenching was purely diffusional.

Control experiments were carried out to explore the roles of Ni and Cr in the catalytic cycle. First, a reaction was performed only in the presence of alkenyl triflate **2s**, NiBr₂, ligand L₃, and HE (Figure 3a). After irradiation with 400 nm LED light for 16 h, dimer **8** was obtained in 12% yield. This result confirmed that




Figure 3. Control experiments to explore the roles of Ni and Cr (for details, see SI S6).

Ni²⁺ could be reduced by HE* and the low-valent Ni species could undergo oxidative addition with **2s**. The resulting **7** should be the key intermediate for the formation of dimer **8** in the catalytic cycle. Furthermore, intermediate **7** should be stable to proton/acid and can last for a long time in the reaction system, because most of **2s** was recovered (81%). The second control experiment was the same as the first one but in the presence of CrCl₂ (Figure 3b). In this case, reduced product **10** was obtained in 73% yield. This result indicated that the transmetalation step that converted **7** to **9** should be fast. Intermediate **9** is highly sensitive to the proton and hydrolyzes rapidly to produce **10**.

Based on these results, we tentatively propose the following catalytic cycle (Figure 4). First, Cr³⁺ is reduced to Cr²⁺ by




Figure 4. Proposed catalytic mechanism.

excited HE *via* a SET process. Ni²⁺ could be reduced to Ni⁰ either by Cr²⁺ or by HE*. The resulting HE* may further participate in electron transfer events by Cr³⁺ and eventually produce pyH⁺. Oxidative addition of alkenyl triflate to low-valent Ni produces intermediate vinyl-Ni **11** which is stable to proton hydrolysis. Transmetalation of **11** with Cr produces nucleophilic vinyl-Cr **12** that is subsequently coupled with carbonyls. After hydrolysis by pyH⁺, the allylic alcohols are obtained and free Cr³⁺ is released.

In summary, the first photochemical NHK coupling has been developed. The additive-free conditions took advantage of the photoreduction ability of HE, bypassing the need for exogenous photosensitizers or metal reductants. The synthetic application was demonstrated through the late-stage derivatization of drug molecules and bioactive skeletons. The unique feature of photochemical NHK is that easily SET-reduced electron-poor substrates, such as pentafluorobenzaldehyde, are accommodated well and produce desired allylic alcohols when classical conditions failed.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.2c00877>.

Experimental details, characterization data of compounds, NMR spectra. (PDF)

AUTHOR INFORMATION

Corresponding Author

Lei Shi – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China; School of Chemistry and Chemical Engineering, Henan Normal University, 453007 Xinxiang, China; orcid.org/0000-0002-2644-1168; Email: shilei17@dlut.edu.cn

Authors

Yonghong Liu – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China

Shuangjie Lin – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China

Dandan Zhang – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China

Bingkun Song – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China

Yunhe Jin – State Key Laboratory of Fine Chemicals, Zhang Dayu School of Chemistry, Dalian University of Technology, 116024 Dalian, China; orcid.org/0000-0003-0626-4587

Erjun Hao – School of Chemistry and Chemical Engineering, Henan Normal University, 453007 Xinxiang, China

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.orglett.2c00877>

Author Contributions

[§]Y.L. and S.L. contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This project is supported by the National Natural Science Foundation of China (2217010858), Open Research Fund of School of Chemistry and Chemical Engineering, Henan Normal University (2020YB03), the Key Scientific Research Projects of Higher Education Institutions in Henan Province (21A150002). We thank WATTCAS CHEM-TECH Co., Ltd. for kindly providing the photoreactor.

REFERENCES

- (1) Zeng, X. Recent Advances in Chromium-Catalyzed Organic Transformations. *Synlett* **2020**, 31 (3), 205–210.
- (2) Cong, X.; Zeng, X. Mechanistic Diversity of Low-Valent Chromium Catalysis: Cross-Coupling and Hydrofunctionalization. *Acc. Chem. Res.* **2021**, 54 (8), 2014–2026.
- (3) Tian, Q.; Zhang, G. Recent Advances in the Asymmetric Nozaki–Hiyama–Kishi Reaction. *Synthesis* **2016**, 48 (23), 4038–4049.
- (4) Gil, A.; Albericio, F.; Álvarez, M. Role of the Nozaki–Hiyama–Takai–Kishi Reaction in the Synthesis of Natural Products. *Chem. Rev.* **2017**, 117 (12), 8420–8446.
- (5) Fürstner, A. Carbon–Carbon Bond Formations Involving Organochromium(III) Reagents. *Chem. Rev.* **1999**, 99 (4), 991–1046.
- (6) Jin, H.; Uenishi, J.; Christ, W. J.; Kishi, Y. Catalytic effect of nickel(II) chloride and palladium(II) acetate on chromium(II)-mediated coupling reaction of iodo olefins with aldehydes. *J. Am. Chem. Soc.* **1986**, 108 (18), 5644–5646.

- (7) Takai, K.; Tagashira, M.; Kuroda, T.; Oshima, K.; Utimoto, K.; Nozaki, H. Reactions of alkenylchromium reagents prepared from alkenyl trifluoromethanesulfonates (triflates) with chromium(II) chloride under nickel catalysis. *J. Am. Chem. Soc.* **1986**, 108 (19), 6048–6050.

- (8) Okude, Y.; Hirano, S.; Hiyama, T.; Nozaki, H. Grignard-type carbonyl addition of allyl halides by means of chromous salt. A chemospecific synthesis of homoallyl alcohols. *J. Am. Chem. Soc.* **1977**, 99 (9), 3179–3181.

- (9) Cintas, P. Addition of Organochromium Compounds to Aldehydes: The Nozaki–Hiyama Reaction. *Synthesis* **1992**, 1992 (03), 248–257.

- (10) Fürstner, A.; Shi, N. Nozaki–Hiyama–Kishi Reactions Catalytic in Chromium. *J. Am. Chem. Soc.* **1996**, 118 (49), 12349–12357.

- (11) Fürstner, A.; Shi, N. A Multicomponent Redox System Accounts for the First Nozaki–Hiyama–Kishi Reactions Catalytic in Chromium. *J. Am. Chem. Soc.* **1996**, 118 (10), 2533–2534.

- (12) Inoue, M.; Suzuki, T.; Nakada, M. Asymmetric Catalysis of Nozaki–Hiyama Allylation and Methallylation with A New Tridentate Bis(oxazolonyl)carbazole Ligand. *J. Am. Chem. Soc.* **2003**, 125 (5), 1140–1141.

- (13) Hargaden, G. C.; Guiry, P. J. The Development of the Asymmetric Nozaki–Hiyama–Kishi Reaction. *Advanced Synthesis & Catalysis* **2007**, 349 (16), 2407–2424.

- (14) Gao, Y.; Hill, D. E.; Hao, W.; McNicholas, B. J.; Vantourout, J. C.; Hadt, R. G.; Reisman, S. E.; Blackmond, D. G.; Baran, P. S. Electrochemical Nozaki–Hiyama–Kishi Coupling: Scope, Applications, and Mechanism. *J. Am. Chem. Soc.* **2021**, 143 (25), 9478–9488.

- (15) Kuroboshi, M. Electrochemical Regeneration of Chromium(II). Alkenylation of Carbonyl Compounds. *Synlett* **1999**, 1999 (1), 69–70.

- (16) Durandetti, M.; Nédélec, J.; Périchon, J. An Electrochemical Coupling of Organic Halide with Aldehydes, Catalytic in Chromium and Nickel Salts. The Nozaki–Hiyama–Kishi Reaction. *Org. Lett.* **2001**, 3 (13), 2073–2076.

- (17) Fiorito, D.; Scaringi, S.; Mazet, C. Transition metal-catalyzed alkene isomerization as an enabling technology in tandem, sequential and domino processes. *Chem. Soc. Rev.* **2021**, 50 (2), 1391–1406.

- (18) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, 116 (17), 10035–10074.

- (19) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, 116 (17), 10075–10166.

- (20) Chen, Y.; Lin, S.; Li, F.; Zhang, X.; Lin, L.; Shi, L. Cp₂Ti^{III}Cl Catalysis in a New Light. *ChemPhotoChem.* **2020**, No. 4, 659–663.

- (21) Schwarz, J. L.; Schäfers, F.; Tlahuext-Aca, A.; Lückemeier, L.; Glorius, F. Diastereoselective Allylation of Aldehydes by Dual Photoredox and Chromium Catalysis. *J. Am. Chem. Soc.* **2018**, 140 (40), 12705–12709.

- (22) Schwarz, J. L.; Kleinmans, R.; Paulisch, T. O.; Glorius, F. 1,2-Amino Alcohols via Cr/Photoredox Dual-Catalyzed Addition of α -Amino Carbanion Equivalents to Carbonyls. *J. Am. Chem. Soc.* **2020**, 142 (5), 2168–2174.

- (23) Tanabe, S.; Mitsunuma, H.; Kanai, M. Catalytic Allylation of Aldehydes Using Unactivated Alkenes. *J. Am. Chem. Soc.* **2020**, 142 (28), 12374–12381.

- (24) Fukuzumi, S.; Hironaka, K.; Tanaka, T. Photoreduction of alkyl halides by an NADH model compound. An electron-transfer chain mechanism. *J. Am. Chem. Soc.* **1983**, 105 (14), 4722–4727.

- (25) Fukuzumi, S.; Inada, O.; Suenobu, T. Mechanisms of Electron-Transfer Oxidation of NADH Analogues and Chemiluminescence. Detection of the Keto and Enol Radical Cations. *J. Am. Chem. Soc.* **2003**, 125 (16), 4808–4816.

- (26) Zhu, D. L.; Wu, Q.; Li, H. Y.; Li, H. X.; Lang, J. P. Hantzsch Ester as a Visible-Light Photoredox Catalyst for Transition-Metal-Free Coupling of Arylhalides and Arylsulfonates. *Chemistry – A European Journal* **2020**, 26 (16), 3484–3488.

- (27) Conrad, K. S.; Manahan, C. C.; Crane, B. R. Photochemistry of flavoprotein light sensors. *Nat. Chem. Biol.* **2014**, *10* (10), 801–809.
- (28) Wang, P. Z.; Chen, J. R.; Xiao, W. J. Hantzsch esters: an emerging versatile class of reagents in photoredox catalyzed organic synthesis. *Org. Biomol. Chem.* **2019**, *17* (29), 6936–6951.
- (29) Buzzetti, L.; Prieto, A.; Roy, S. R.; Melchiorre, P. Radical-Based C–C Bond-Forming Processes Enabled by the Photoexcitation of 4-Alkyl-1,4-dihydropyridines. *Angew. Chem., Int. Ed.* **2017**, *56*, 15039–15043.
- (30) Kammer, L. M.; Badir, S. O.; Hu, R.; Molander, G. A. Photoactive electron donor–acceptor complex platform for Ni-mediated C(sp³)–C(sp²) bond formation. *Chemical Science* **2021**, *12* (15), 5450–5457.
- (31) Lin, S.; Chen, Y.; Yan, H.; Liu, Y.; Sun, Y.; Hao, E.; Shi, C.; Zhang, D.; Zhu, N.; Shi, L. Activation of Chromium Catalysts by Photoexcited Hantzsch Ester for Decarboxylative Allylation of Aldehydes with Butadiene. *Org. Lett.* **2021**, *23* (20), 8077–8081.
- (32) Liu, X.; Li, X.; Chen, Y.; Hu, Y.; Kishi, Y. On Ni Catalysts for Catalytic, Asymmetric Ni/Cr-Mediated Coupling Reactions. *J. Am. Chem. Soc.* **2012**, *134* (14), 6136–6139.
- (33) Jung, J.; Kim, J.; Park, G.; You, Y.; Cho, E. J. Selective Debromination and α -Hydroxylation of α -Bromo Ketones Using Hantzsch Esters as Photoreductants. *Advanced Synthesis & Catalysis* **2016**, *358* (1), 74–80.
- (34) Huang, H. M.; Bellotti, P.; Daniliuc, C. G.; Glorius, F. Radical Carbonyl Propargylation by Dual Catalysis. *Angew. Chem., Int. Ed.* **2021**, *60*, 2464–2471.
- (35) Till, N. A.; Smith, R. T.; MacMillan, D. W. C. Decarboxylative Hydroalkylation of Alkynes. *J. Am. Chem. Soc.* **2018**, *140* (17), 5701–5705.

Recommended by ACS

Unlocking C–H Functionalization at Room Temperature via a Light-Mediated Protodemetalation Reaction

Claire Empel, Rene M. Koenigs, *et al.*

JUNE 26, 2022
ACS CATALYSIS

READ 

Photoinduced C(sp³)-H Chalcogenation of Amide Derivatives and Ethers via Ligand-to-Metal Charge-Transfer

Ben Niu, Sébastien Laulhé, *et al.*

JUNE 21, 2022
ORGANIC LETTERS

READ 

Photocatalyzed Dehydrogenation of Aliphatic N-Heterocycles Releasing Dihydrogen

, Burkhard König, *et al.*

AUGUST 08, 2022
ACS CATALYSIS

READ 

Visible-Light-Promoted Fe(III)-Catalyzed N–H Alkylation of Amides and N-Heterocycles

Hangcheng Ni, Hui Mao, *et al.*

JULY 20, 2022
THE JOURNAL OF ORGANIC CHEMISTRY

READ 

Get More Suggestions >